

# **Doctors please Fax this Referral to 9194-7789**

Your patient will receive an appointment letter in the mail Appointments cannot be made over the phone without this referral

# Respiratory.Bookings@easternhealth.org.au

#### **ALL TESTS BULK BILLED** Please fill all fields, legibly - help us to help you Surname Name Sex Address M UR F Medicare DOB Phone: Clinical Question – suggested tests listed Spirometry/TLCO ± Lung Volumes if normal, Bronchial Challenge □ Cough FI / Asthma? Dyspnoea FI / COPD / Emphysema / OSA / Pre-operative assessment Interstitial lung disease or Fibrosis / Pulmonary Hypertension / Other (Please specify) Spirometry/TLCO ± Lung Volumes ± Cardiopulmonary Exercise Test (CPET) Asthma for review Spirometry/Expired Nitric Oxide (FeNO) Spirometry/Expired Nitric Oxide (FeNO)/Skin Prick Test (Aeroallergens) **Allergies** Oxygen assessment for flight Hypoxic Altitude Simulation Test (HAST) Oxygen Assessment Spirometry/TLCO ± Lung Volumes/6MWT/ABG and O2 Consult MIPS & MEPS\* Other tests: Anatomic Shunt Fraction\* **Tests Ordered** □ Spirometry □ TLCO±Lung Volumes □ Exhaled Nitric Oxide ■ Skin Prick Tests □ Bronchial Challenge ■ MIPS & MEPS ☐ 6MWT □ HAST ■ Anatomic Shunt Fraction □ ABGs □ Oxygen consultation □ CPET

Department of	
Respiratory Medic	ine

**Box Hill Hospital** Monday to Friday

Respiratory Laboratory, 3N, Third Floor, Building B, Nelson Rd

# Maroondah Hospital

Tuesdays and Thursdays Only Davey Drive (near Mt Dandenong Rd) Ringwood East

\* Lung Volumes, Skin Prick Test, HAST only available Box Hill

Other Clinical Data:

Referring Doctor		
Dr	Provider No	
Address		
Phone	Fax	
Signature	Date	
Cinnet up not sequiped if unleaded alestropically		

#### **ROUTINE TESTS AVAILABLE**

- Spirometry Measurement of FEV1 and FVC. Designed to identify obstruction & measure airway reversibility (asthma).
- TLCO Measures the lungs ability to transfer oxygen from the alveoli into the pulmonary capillary blood. Varies with: Hb, thickness of alveolarcapillary membrane (fibrosis), surface area of the alveolar-capillary membrane (emphysema).
- Expired Nitric Oxide (FeNO) Determines the presence of current eosinophilic airway inflammation.
- Lung Volumes (Plethysmography) Measures lung size (eg. TLC) to confirm restriction.
- MIPS & MEPS Maximal Inspiratory and Expiratory Pressures: Quantifies respiratory muscle strength.
- Bronchial Challenge Measures bronchial hyper reactivity, to assist in the diagnosis of current asthma.
- Skin Prick Test (aeroallergens only) Determines the presence of skin sensitizing antibodies (IgE) to aid in the diagnosis of allergic disease.

test. Don't Use creams, moisturiser etc on your arms on the day of the test.

Copy of results sent to:

Follow up Doctors appointment:

Tel: 9895 3437

**Date** Time

### SPECIALISED TESTS AVAILABLE

- 6MWT (6-minute walk) including ABG, determines need for domiciliary oxygen at rest and during exercise.
- HAST (Hypoxic Altitude Simulation Test) Assess patient's oxygen requirements at a simulated altitude for air travel.
- Shunt Fraction Measurement of the anatomic shunt.
- Cardiopulmonary exercise test Assess the nature and severity of cardiovascular and pulmonary disease.

# **CONTRAINDICATIONS**

- Abdominal, thoracic, neuro or ocular surgery past 6/52
- Pneumothorax in past 6 weeks.
- Haemoptysis of unknown origin.
- Open pulmonary TB
- Unstable cardiovascular status.
- Thoracic or cerebral aneurysm.
- Angiogram or Transbronchial biopsy in past 24 hours.
- Intercostal catheters in situ.
- Recent myocardial infarction or pulmonary embolus.

Please STOP: Ventolin, Airomir, Asmol, Bricanyl at least 4 hours prior to test, unless you really feel you need it, Atrovent, Serevent, Seretide, Oxis, Symbicort, Flutiform, Bretaris, Brimica, DuoResp, Fostair, Fluticasone+Salmetarol Cipla at least 12 hours prior. Spiriva, Braltus, Onbrez, Seebri, Spiolto, Ultibro, Anoro, Incruse, Breo, Singulair, Spiolto, Trelegy at least 24 hours prior. You may continue: Flixotide, Qvar, Alvesco, Arnuity, Pulmicort, Flutacosone Cipla. No smoking for at least 6 hours prior to test.

Bronchial Challenge: STOP all of the above, including Ventolin, Airomir, Asmol, Bricanyl at least 8 hours prior to test, Braltus, Spiriva, Onbrez, Bretaris, Seebri, Spiolto, Ultibro, Anoro, Brimica, Incruse at least 72 hours (3 days), no anti-histamines for 3 days. Also STOP Flixotide, Qvar, Alvesco, Arnuity, Seretide, Flutiform, Fostair, Breo, Symbicort, Pulmicort, DuoResp, Fluticasone+Salmetarol Cipla, Flutacosone Cipla, Trelegy for 6 weeks. Atrovent for 12 hours, Serevent, Oxis, Theophylline for 24 hours Singulair for 4 days. DO NOT have coffee, tea, coke, chocolate (anything with caffeine) and Vitamin C

supplements from midnight prior to the test. No vigorous exercise on the day of the test. Skin Prick Tests: STOP Hay fever and allergy medications, including Telfast, Fexotabs, Claratyne, Clarinase, Zadine, Zyrtec, Polaramine, for 3 days before the

FeNO breath test: For at least one hour prior to the test - Don't eat or drink (including NO water), don't perform strenuous exercise and don't smoke. DON'T eat any processed meats on the day eg bacon, salami, sausages, ham etc.